Facilitating Recovery After Stroke: Traditional Methods Meet Technologically Innovative Interventions

University of Michigan Health System
Robert Ferguson, OTR/L
Douglas Rakoski MA, OTR/L, ATP

Evidence-Based Reviews and Analyses

Barclay-Goddard RE, Stevenson TJ, Poluha W, Thalman L. Mental practice for treating upper extremity deficits in individuals with hemiparesis after stroke. Cochrane Database Syst Rev 2011; 5:CD005950.

Bolton D, Cauraugh JH, Hausenblas HA. Electromyogram-triggered neuromuscular stimulation and stroke motor recovery of arm/hand functions: a meta-analysis. J Neurol Sci 2004;223:121–7.

Bonaiuti D, Rebasti L, Sioli P. The constraint induced movement therapy: a systematic review of randomised controlled trials on the adult stroke patients. Eura Medicophys 2007;43:139-146

Braun SM, Beurskens AJ, Borm PJ, Schack T, Wade DT. The effects of mental practice in stroke rehabilitation: a systematic review. Arch Phys Med Rehabil 2006;87:842–52.

Cauraugh JH, Lodha N, Naik SK, Summers JJ. Bilateral movement training and stroke motor recovery progress: a structured review and meta-analysis. Hum Mov Sci 2010; 29(5):853-870.

Cooke EV, Mares K, Clark A, Tallis RC, Pomeroy VM. The effects of increased dose of exercise-based therapies to enhance motor recovery after stroke: a systematic review and meta-analysis. BMC Med 2010; 8:60.

Corbetta D, Sirtori V, Moja L, Gatti R. Constraint-induced movement therapy in stroke patients: systematic review and metaanalysis. Eur J Phys Rehabil Med 2010;46(4):537-544.

Coupar F, Pollock A, van WF, Morris J, Langhorne P. Simultaneous bilateral training for improving arm function after stroke. Cochrane Database Syst Rev 2010;4:CD006432.

de Kroon JR, van der Lee JH, IJzerman MJ, Lankhorst GJ. Therapeutic electrical stimulation to improve motor control and functional abilities of the upper extremity after stroke: a systematic review. Clin Rehabil 2002,16(4):350-360.

Hakkenes S, Keating JL. Constraint-Induced movement therapy following stroke systematic review of randomised controlled trials. Australian Journal of Physiotherapy 2005;51:221-231.

Henderson A, Korner-Bitensky N, Levin M. Virtual reality in stroke rehabilitation: a systematic review of its effectiveness for upper limb motor recovery. Top Stroke Rehabil 2007;14:52-61.

rferg@med.umich.edu
drakoski@med.umich.edu

Hiraoka K. Rehabilitation effort to improve upper extremity function in post-stroke patients:A meta-analysis. J Phys Ther Sci 2005;13:5-9.

Kwakkel G, Kollen BJ, Krebs HI. Effects of robot-assisted therapy on upper limb recovery after stroke: a systematic review. Neurorehabil Neural Repair 2008;22:111-121.

Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. Lancet Neurol 2009;8:741–754.

Latimer CP, Keeling J, Lin B, Henderson M, Hale LA. The impact of bilateral therapy on upper limb function after chronic stroke: a systematic review. Disabil Rehabil 2010;32:1221-1231.

Laver KE, George S, Thomas S, Deutsch JE, Crotty M. Virtual reality for stroke rehabilitation. Cochrane Database Syst Rev 2011; 9:CD008349.

Masiero S, Armani M, Rosati G. Upper-limb robot-assisted therapy in rehabilitation of acute stroke patients: Focused review and results of new randomized controlled trial. J Rehabil Res Dev 2011; 48(4):355-366.

Meilink A, Hemmen B, Seelen HA, Kwakkel G. Impact of EMG-triggered neuromuscular stimulation of the wrist and finger extensors of the paretic hand after stroke: a systematic review of the literature. Clin Rehabil 2008;22:291-305.

Nilsen DM, Gillen G, Gordon AM. Use of mental practice to improve upper-limb recovery after stroke: a systematic review. Am J Occup Ther 2010; 64(5):695-708.

Paci M. Physiotherapy based on the Bobath concept for adults with poststroke hemiplegia: a review of effectiveness studies. J Rehabil Med 2003;35:2–7.

Pomeroy VM, King L, Pollock A, Baily-Hallam A,Langhorne P. Electrostimulation for promoting recovery of movement or functional ability after stroke. Cochrane Database Syst Rev 2006;(2):CD003241.

Prange GB, Jannink MJ, Groothuis-Oudshoorn CG et al. Systematic review of the effect of robot-aided therapy on recovery of the hemiparetic arm after stroke. J Rehabil Res Dev. 2006;43:171-184.

Richards LG, Stewart KC, Woodbury ML, Senesac C, Cauraugh JH. Movement-dependent stroke recovery: a systematic review and meta-analysis of TMS and fMRI evidence. Neuropsychologia 2008;46:3–11.

Saposnik G, Levin M. Virtual Reality in Stroke Rehabilitation: A Meta-Analysis and Implications for Clinicians. Stroke 2011; 42(5):1380-1386.

Shi YX, Tian JH, Yang KH, Zhao Y. Modified constraint-induced movement therapy versus traditional rehabilitation in patients with upper-extremity dysfunction after stroke: a systematic review and meta-analysis. Arch Phys Med Rehabil 2011; 92(6):972-982.

Sirtori V, Corbetta D, Moja L, Gatti R. Constraint-induced movement therapy for upper extremities in stroke patients. Cochrane Database Syst Rev 2009;CD004433.

rferg@med.umich.edu
drakoski@med.umich.edu

Steultjens EM, Dekker J, Bouter LM, van de Nes JC, Cup EH, van den Ende CH. Occupational therapy for stroke patients: a systematic review. Stroke 2003;34:676-87.

Stewart KC, Cauraugh JH, Summers JJ. Bilateral movement training and stroke rehabilitation: a systematic review and metaanalysis. J Neurol Sci 2006;244:89-95.

Subramanian SK, Massie CL, Malcolm MP, Levin MF. Does provision of extrinsic feedback result in improved motor learning in the upper limb poststroke? A systematic review of the evidence. Neurorehabil Neural Repair 2010;24:113-124.

Teasell RW, Foley NC, Sanjit KB, Speechley MR. An evidence-based review of stroke rehabilitation. Top Stroke Rehabil. 2003; 10(1): 29-58

Timmermans AA, Spooren AI, Kingma H, Seelen HA. Influence of task-oriented training content on skilled arm-hand performance in stroke: a systematic review. Neurorehabil Neural Repair 2010; 24(9):858-870.

Tyson SF, Kent RM. The effect of upper limb orthotics after stroke: a systematic review. NeuroRehabilitation 2011; 28(1):29-36.

Van Dijk H, Jannink M, Hermens H. Effect of augmented feedback on motor function of the affected upper extremity in rehabilitation patients: a systematic review of randomized controlled trials. J Rehabil Med 2005;37:202–11.

Woldag H, Hummelsheim H. Evidence-based physiotherapeutic concepts for improving arm and hand function in stroke patients: a review. J Neurol 2002;249(5):518-28.

Zimmermann-Schlatter A, Schuster C, Puhan MA, Siekierka E, Steurer J. Efficacy of motor imagery in post-stroke rehabilitation: a systematic review. J Neuroeng Rehabil 2008;5:8.

rferg@med.umich.edu
drakoski@med.umich.edu

Bibliography

Alon G, Levitt AF, McCarthy PA. Functional electrical stimulation enhancement of upper extremity functional recovery during stroke rehabilitation: a pilot study. Neurorehabil Neural Repair 2007;21:207–15.

Altschuler EL, Wisdom S, Stone L, Foster C, Galasko D. Rehabilitation of hemiparesis after stroke with a mirror. Lancet 1999;353:2035–6.

Altschuler EL, Wisdom SB, Stone L, Foster C, Galasko D, Llewellyn DM, Ramachandran VS. Rehabilitation of hemiparesis after stroke with a mirror. Lancet 1999;353:2035-2036.

Barker RN, Brauer SG. Upper limb recovery after stroke: the stroke survivors' perspective. Disabil Rehabil 2005;27:1213–1223.

Barzel A, Liepert J, Haevernick K, Eisele M, Ketels G, Rijntjes M, van den BH. Comparison of two types of Constraint- Induced Movement Therapy in chronic stroke patients: A pilot study. Restor Neurol Neurosci 2009;27:673-680.

Blennerhassett J, Dite W. Additional task related practice improves mobility and upper limb function early after stroke: a randomised controlled trial. Aust J Physiother 2004;50:219-224.

Blennerhassett J, Dite W. Additional task-related practice improves mobility and upper limb function early after stroke: a randomized controlled trial. Aust J Physiother 2004;50:219–24.

Boake C, Noser EA, Ro T, Baraniuk S, Gaber M, Johnson R, et al. Constraint-induced movement therapy during early stroke rehabilitation. Neurorehabil Neural Repair 2007;21:14–24.

Boake C, Noser EA, Ro T, Baraniuk S, Gaber M, Johnson R, Salmeron ET, Tran TM, Lai JM, Taub E, Moye LA, Grotta JC, Levin HS. Constraint-Induced Movement Therapy During Early Stroke Rehabilitation. Neurorehabil Neurol Repair 2007;21:14-24.

Bobath, B. (Ed.). Adult Hemiplegia: evaluation and treatment. London: Heinemann Medical Books; 1990.

Bovend'Eerdt TJ, Dawes H, Sackley C, Izadi H, Wade DT. An integrated motor imagery program to improve functional task performance in neurorehabilitation: a singleblind randomized controlled trial. Arch Phys Med Rehabil 2010;91:939-946.

Brogardh C, Flansbjer UB, Lexell J. What is the long-term benefit of constraint-induced movement therapy? A four-year follow-up. Clin Rehabil 2009a;23:418-423.

Brogardh C, Vestling M, Sjolund BH. Shortened constraint-induced movement therapy in subacute stroke - no effect of using a restraint: a randomized controlled study with independent observers. J Rehabil Med 2009;41:231-236.

rferg@med.umich.edu
drakoski@med.umich.edu

Brown J, Lutsep H, Weinand M, Cramer S. Motor cortex stimulation for the enhancement of recovery from stroke: a prospective, multicenter safety study. Neurosurgery 2006;58:464–73.

Brunnstrom S. Movement Therapy in Hemiplegia. New York: Harper & Row; 1970.

Burgar CG, Lum PS, Scremin AM, Garber SL, Van der Loos HF, Kenney D et al. Robot assisted upper-limb therapy in acute rehabilitation setting following stroke: Department of Veterans Affairs multisite clinical trial. J Rehabil Res Dev 2011; 48(4):445-458.

Butefisch C, Hummelsheim H, Denzler P, Mauritz KH. Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. J Neurol Sci 1995;130(1):59-68.

Butler AJ, Page SJ. Mental practice with motor imagery: evidence for motor recovery and cortical reorganization after stroke. Arch Phys Med Rehabil 2006;87:S2-11

Byl N, Roderick J, Mohamed O, Hanny M, Kotler J, Smith A, Tang M, Abrams G. Effectiveness of sensory and motor rehabilitation of the upper limb following the principles of neuroplasticity: patients stable post-stroke. Neurorehabil Neural Repair 2003;17(3):176-91.

Byl NN, Pitsch EA, Abrams GM. Functional outcomes can vary by dose: learning-based sensorimotor training for patients stable post-stroke. Neurorehabil Neural Repair 2008;22:494-04.

Cardoso de Oliveira S. The neuronal basis of bimanual coordination: recent neurophysiological evidence and functional models. Acta Psychologica 2002;110:139–59.

Carr J.H and Shepherd R.B. A Motor Relearning Programme for Stroke 2nd ed. London: William Heinemann Medical Books (1987).

Carr JH, Shepherd RB. Stroke rehabilitation: guidelines for exercises and training to optimize motor skill. Oxford: Butterworth-Heinemann; 2002.

Castel-Lacanal E, Gerdelat-Mas A, Marque P, Loubinoux I, Simonetta- Moreau M. Induction of cortical plastic changes in wrist muscles by paired associative stimulation in healthy subjects and post-stroke patients. Exp Brain Res 2007;180:113–22.

Cauraugh J, Light K, Kim SB, Thigpen M, Behrman A. Chronic motor dysfunction after stroke: recovering wrist and finger extension by electromyography-triggered neuromuscular stimulation. Stroke 2000;31(6):1360-4.

Cauraugh JH, Coombes SA, Lodha N, Naik SK, Summers JJ. Upper extremity improvements in chronic stroke: coupled bilateral load training. Restor Neurol Neurosci 2009;27:17-25.

Cauraugh JH, Kim SB . Two coupled motor recovery protocols are better than one: electromyogram-triggered neuromuscular stimulation and bilateral movements. Stroke 2002; 33(6):1589-1594.

rferg@med.umich.edu
drakoski@med.umich.edu

Cauraugh JH, Kim SB. Stroke motor recovery: active neuromuscular stimulation and repetitive practice schedules. J Neurol Neurosurg Psychiatry 2003;74(11):1562-6.

Cauraugh JH, Kim SB. Chronic stroke motor recovery: duration of active neuromuscular stimulation. J Neurol Sci 2003;215:13–9.

Cauraugh JH, Kim SB, Duley A. Coupled bilateral movements and active neuromuscular stimulation: intralimb transfer evidence during bimanual aiming. Neurosci Lett 2005;382:39–44.

Chan MK, Tong RK, Chung KY. Bilateral upper limb training with functional electric stimulation in patients with chronic stroke. Neurorehabil Neural Repair 2009;23:357-365.

Church C, Price C, Pandyan AD, Huntley S, Curless R, Rodgers H. Randomized controlled trial to evaluate the effect of surface neuromuscular electrical stimulation to the shoulder after acute stroke. Stroke 2006;37:2995–3001.

Cirstea MC, Levin ML. Improvement of arm movement patterns and endpoint control depends on type of feedback during practice in stroke survivors. Neurorehabil Neural Repair 2007;21:398–411.

Classen J, Liepert J, Wise SP, Hallett M, Cohen LG. Rapid plasticity of human cortical movement representation induced by practice. J Neurophysiol 1998;79(2):1117- 23.

Daly JJ, Hogan N, Perepezko EM, Krebs HI, Rogers JM, Goyal KS, et al. Response to upper-limb robotics and functional neuromuscular stimulation following stroke. J Rehabil Res Dev 2005;42:723–36.

Davis J, Task Selection and Enriched Environments: A functional upper extremity training program for stroke survivors. Topics Stroke Rehabil, 2006; 13(3): 1-11.

de Kroon JR, Ijzerman MJ, Chae J. Relation between stimulation characteristics and clinical outcome in studies using electrical stimulation to improve motor control of the upper extremity in stroke. J Rehabil Med 2005;37:65–74.

de Kroon JR, IJzerman MJ, Lankhorst GJ, Zilvold G. Electrical stimulation of the upper limb in stroke: stimulation of the extensors of the hand vs. alternate stimulation of flexors and extensors. Am J Phys Med Rehabil 2004;83(8):592-600.

de Kroon, Jr., IJzerman MJ. Electrical stimulation of the upper extremity in stroke: cyclic versus EMG-triggered stimulation. Clin Rehabil 2008;22:690-697.

Dean CM, Shepherd RB. Task-related training improves performance of seated reaching tasks after stroke. A randomized controlled trial. Stroke 1997;28:722-728.

Desrosiers J, Bourbonnais D, Corriveau H, Gosselin S, Bravo G. Effectiveness of unilateral and symmetrical bilateral task training for arm during the sub-acute phase after stroke: a randomized controlled trial. Clin Rehab 2005;19:581–93.

rferg@med.umich.edu
drakoski@med.umich.edu

Desrosiers J, Bourbonnais D, Corriveau H, Gosselin S, Bravo G. Effectiveness of unilateral and symmetrical bilateral task training for arm during the subacute phase after stroke: a randomized controlled trial. Clin Rehabil 2005;19:581-593.

Dijkerman H, Letswaart M, Johnston M, MacWalter RS. Does motor imagery training improve hand function in chronic stroke patients? A pilot study. Clin Rehab 2003;18:538–49.

Diserens K, Perret N, Chatelain S, Bashir S, Ruegg D, Vuadens P, Vingerhoets F. The effect of repetitive arm cycling on post stroke spasticity and motor control Repetitive arm cycling and spasticity. J Neurol Sci 2007;253:18-24.

Dohle C, Pullen J, Nakaten A, Kust J, Rietz C, Karbe H. Mirror therapy promotes recovery from severe hemiparesis: a randomized controlled trial. Neurorehabil Neural Repair 2009;23:209-217.

Donaldson C, Tallis R, Miller S, Sunderland A, Lemon R, Pomeroy V. Effects of conventional physical therapy and functional strength training on upper limb motor recovery after stroke: a randomized phase II study. Neurorehabil Neural Repair 2009;23:389-397.

Dromerick AW, Lang CE, Birkenmeier RL, Wagner JM, Miller JP, Videen TO, Powers WJ, Wolf SL, Edwards DF. Very early constraint-induced movement during stroke rehabilitation (VECTORS): A single-center RCT. Neurology 2009;73:195-201.

Ertelt D, Small S, Solodkin A, Dettmers C, McNamara A, Binkofski F. Action Observation has positive impact on rehabilitation of motor deficits after stroke. Neuroimage 2007;36:164-173.

Fasoli SE, Krebs HI, Stein J, Frontera W, Hogan N. Effects of robotic therapy on motor impairment and recovery in chronic stroke. Arch Phys Med Rehabil 2003;84:477–82.

Fasoli SE, Krebs HI, Stein J, Frontera WR, Hughes R, Hogan N. Robotic therapy for chronic motor impairments after stroke: Follow-up results. Arch Phys Med Rehabil 2004; 85(7):1106- 111.

Feys H, De Weerdt W, Werbeke G, Steck G, Capiau C, Kiekens C, et al. Early and repetitive stimulation of the arm can substantially improve the long-term outcome after stroke: a 5-year follow up study of a randomized trial. Stroke 2004;35:924–9.

Finley MA, Fasoli SE, Dipietro L, Ohlhoff J, MacClellan L, Meister C, Whitall J, Macko R, Bever CT, Krebs HI, Hogan N. Short-duration robotic therapy in stroke patients with severe upper-limb motor impairment. J Rehabil Res Dev 2005;42:683-692.

Fischer HC, Stubblefield K, Kline T, Luo T, Kenyon RV, Kamper DG. Hand rehabilitation following stroke: a pilot study of assisted finger extension training in a virtual environment. Top Stroke Rehabil 2007;14:1-12.

Floel A, Nagorsen U, Werhahn K, Ravindran S. Influence of somatosensory input on motor function in patients with chronic stroke. Ann Neurol 2004;56:206–12.

rferg@med.umich.edu
drakoski@med.umich.edu

Fregni F, Boggio PS, Valle AC, Rocha RR, Duarte J, Ferreira MJL, et al. A sham-controlled trial of a 5-day course of repetitive transcranial magnetic stimulation of the unaffected hemisphere in stroke patients. Stroke 2006;37:2115–22.

Fritz S, Chiu Y, Malcolm M, Patterson T, Light K. Feasibility of electromyography-triggered neuromuscular stimulation as an adjunct to constraint-induced movement therapy. Phys Ther 2005;85: 428–42.

Gabr U, Levine P, Page SJ. Home-based electromyography-triggered stimulation in chronic stroke. Clin Rehab 2005;19:737–45.

Gabr U, Levine P, Page SJ. Home-based electromyography-triggered stimulation in chronic stroke. Clin Rehabil 2005; 19(7):737-745.

Gauthier LV, Taub E, Perkins C, Ortmann M, Mark VW, Uswatte G. Remodeling the brain. Plastic changes produced by different motor therapies after stroke. Stroke 2008;39:1520–5.

Gilmore PE, Spaulding SJ. Motor learning and the use of videotape feedback after stroke. Top Stroke Rehabil 2007;14:28-36.

Gritsenko V, Prochazka A. A functional electric stimulation-assisted exercise therapy system for hemiplegic hand function. Arch Phys Med Rehabil 2004;85:881–5.

Hafsteinsdóttir TB, Algra A, Kappelle LJ, Grypdonck. Neurodevelopmental treatment after stroke: a comparative study. J Neurol Neurosurg Psychiatry 2005; 76:788-792.

Hafsteinsdottir TB, Kappelle J, Grypdonck MH, Algra A. Effects of Bobath-based therapy on depression, shoulder pain and health-related quality of life in patients after stroke. J Rehabil Med 2007;39:627-632

Hallett M. Transcranial magnetic stimulation and the human brain. Nature 2000;406:147–50.

Hammer A, Lindmark B. Is forced use of the paretic upper limb beneficial? A randomized pilot study during sub-acute post-stroke recovery. Clin Rehabil 2009a;23:424-433.

Hammer AM, Lindmark B. Effects of forced use on arm function in the subacute phase after stroke: a randomized, clinical pilot study. Phys Ther 2009b;89:526-539.

Hamzei F, Liepert J, Dettmers C, Weiller C, Rijntjes M. Two different reorganization patterns after rehabilitative therapy: an exploratory study with fMRI and TMS. Neuroimage 2006;31:710–20.

Hara Y, Ogawa S, Muraoka Y. Hybrid power-assisted functional electrical stimulation to improve hemiparetic upper-extremity function. AmJ Phys Med Rehabil 2006;85:977–85.

Hara Y, Ogawa S, Tsujiuchi K, Muraoka Y. A home-based rehabilitation program for the hemiplegic upper extremity by powerassisted functional electrical stimulation. Disabil Rehabil 2008;30:296-304.

rferg@med.umich.edu
drakoski@med.umich.edu

Harris JE, Eng JJ. Paretic upper-limb strength best explains arm activity in people with stroke. Phys Ther 2007;87:88–97.

Harris-Love M, Mc Combe S, Whitall J. Exploiting interlimb coupling to improve paretic arm reaching performance in people with chronic stroke. Arch Phys Med Rehabil 2005;86:2131–7.

Hayner K, Gibson G, Giles GM. Comparison of constraint-induced movement therapy and bilateral treatment of equal intensity in people with chronic upper-extremity dysfunction after cerebrovascular accident. Am J Occup Ther 2010; 64(4):528-539.

Hemmen B, Seelen HA. Effects of movement imagery and electromyography-triggered feedback on arm hand function in stroke patients in the subacute phase. Clin Rehabil 2007;21:587-594.

Hesse S, Schulte-Tigges G, Konrad M, Bardeleben A, Werner C. Robot-assisted arm trainer for the passive and active practice of bilateral forearm and wrist movements in hemiparetic subjects. Arch Phys Med Rehabil 2003;84:915-20.

Hesse S, Waldner A, Mehrholz J, Tomelleri C, Pohl M, Werner C. Combined transcranial direct current stimulation and robot-assisted arm training in subacute stroke patients: An exploratory, randomized multicenter trial. Neurorehabil Neural Repair 2011(e-publ).

Hesse S, Werner C, Pohl M, Mehrholz J, Puzich U, Krebs HI. Mechanical arm trainer for the treatment of the severely affected arm after a stroke: a single-blinded randomized trial in two centers. Am J Phys Med Rehabil 2008;87:779-788.

Hesse S, Werner C, Pohl M. Computerized arm training improves the motor control of the severely affected arm after stroke: a single-blinded randomized trial in two centers. Stroke 2005;36:1960–6.

Higgins J, Salbach NM, Wood-Dauphinee S, Richards CL, Coˆte´ R, Mayo NE. The effect of task-oriented intervention on arm function in people with stroke: a randomized controlled trial. Clin Rehab 2006; 20:296–310.

Holden M, Dyar T. Virtual environment training: a new tool for neurorehabilitation. Neurol Rep. 2002;26:62-71.

Housman SJ, Scott KM, Reinkensmeyer DJ. A randomized controlled trial of gravity supported, computer-enhanced arm exercise for individuals with severe hemiparesis. Neurorehabil Neural Repair 2009;23:505-514.

Hsieh Y, Wu C, Lin K, Yao G, Wu K, Chang Y. Dose–Response Relationship of Robot-Assisted Stroke Motor Rehabilitation: The Impact of Initial Motor Status. Stroke. 2012; 43: 2729-2734

Hsieh YW, Wu CY, Liao WW, Lin KC, Wu KY, Lee CY. Effects of treatment intensity in upper limb robot-assisted therapy for chronic stroke: a pilot randomized controlled trial. Neurorehabil Neural Repair 2011; 25(6):503-511.

rferg@med.umich.edu
drakoski@med.umich.edu

Hsu SS, Hu MH, Wang YH, Yip PK, Chiu JW, Hsieh CL. Dose-response relation between neuromuscular electrical stimulation and upper-extremity function in patients with stroke. Stroke 2010;41:821-824.

Hu XL, Tong KY, Song R, Zheng XJ, Leung WW. A comparison between electromyography-driven robot and passive motion device on wrist rehabilitation for chronic stroke. Neurorehabil Neural Repair 2009;23:837-846.

Hummel F, Celnik P, Giraux P, Floel A,Wu WH, Gerloff C, et al. Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke. Brain 2005;128:490–9.

Hunter SM, Hammett L, Ball S, Smith N, Anderson C, Clark A et al. Dose-response study of mobilisation and tactile stimulation therapy for the upper extremity early after stroke: a phase I trial. Neurorehabil Neural Repair 2011; 25(4):314-322.

Ietswaart M, Johnston M, Dijkerman HC, Joice S, Scott CL, MacWalter RS et al. Mental practice with motor imagery in stroke recovery: randomized controlled trial of efficacy. Brain 2011; 134(Pt 5):1373-1386.

Jang SH, You SH, Hallett M, Cho YW, Park CM, Cho SH. Cortical reorganization and associated functional motor recovery after virtual reality in patients with chronic stroke: an experimenter-blind preliminary study. Arch Phys Med Rehabil 2005;86:2218–23.

Jeannerod M, Frak V. Mental imaging of motor activity in humans. Curr Opin Neurobiol 1999;9:735–9.

Kahn LE, Zygman ML, Rymer WZ, Reinkensmeyer DJ. Robot-assisted reaching exercise promotes arm movement recovery in chronic hemiparetic stroke: a randomized controlled pilot study. J Neuro Eng Rehab 2005;3:12.

Kalra L, Ratan R. Recent advances in stroke rehabilitation. Stroke 2006;38:235–237.

Kim YH, You SH, Ko MH. Repetitive transcranial magnetic stimulation-induced corticomotor excitability and associated motor skill acquisition in chronic stroke. Stroke 2006;37:1471–6.

Kimberley TJ, Lewis SM, Auerbach EJ, Dorsey LL, Lojovich JM, Carey JR. Electrical stimulation driving functional improvements and cortical changes in subjects with stroke. Exp Brain Res 2004;154:450-460.

Knutson JS, Harley MY, Hisel TZ, Hogan SD, Maloney MM, Chae J. Contralaterally controlled functional electrical stimulation for upper extremity hemiplegia: An early-phase randomized clinical trial in subacute stroke patients. Neurorehabil Neural Repair 2011;26(3):239-46

Kowalczewski J, Gritsenko V, Ashworth N, Ellaway P, Prochazka A. Upper-extremity functional electric stimulation-assisted exercises on a workstation in the subacute phase of stroke recovery. Arch Phys Med Rehabil 2007;88:833-839.

Kwakkel G, Wagenaar R, Twisk J, Lankhorst G, Koetsier J. Intensity of leg and arm training after primary middle cerebral artery stroke: a randomised trial. Lancet 1999;354:191–6.

rferg@med.umich.edu
drakoski@med.umich.edu

Kwakkel G, Wagenaar RC, Twisk JW, Lankhorst GJ, Koetsier JC. Intensity of leg and arm training after primary middle-cerebral-artery stroke: a randomised trial. Lancet 1999;354(9174):191-196.

Lam YS, Man DWK, Tam SF, Weiss PL. Virtual reality training for stroke rehabilitation. Neurorehabilitation 2006;21:245-253.

Langhammer B, Stanghelle JK. Bobath or motor relearning programme? A follow-up one and four years post stroke. Clin Rehabil 2003;17(7):731-734.

Langhammer B, Stanghelle JK. Can physiotherapy after stroke based on the Bobath concept result in improved quality of movement compared to the motor relearning programme. Physiother. Res. Int. 16 (2011) 69–80.

Levy R, Ruland S, Weinand M, Lowry D, Dafer R. Cortical stimulation for rehabilitation of patients with hemiparetic stroke: a multicenter feasibility study of safety and efficacy. J Neurosurg 2008;108:707–14.

Lewis G, ByblowW. Neurophysiological and behavioural adaptations to bilateral training intervention in individuals following stroke. Clin Rehab 2004;18:48–59.

Liao WW, Wu CY, Hsieh YW, Lin KC, Chang WY. Effects of robot-assisted upper limb rehabilitation on daily function and real-world arm activity in patients with chronic stroke: a randomized controlled trial. Clin Rehabil 2012; 26(2): 111-120

Liepert J, Bauder H, Wolfgang HR. Treatment induced cortical reorganization after stroke in humans. Stroke 2000;31:1210–6.

Liepert J, Zittel S,Weiller C. Improvement of dexterity by single session low-frequency repetitive transcranial magnetic stimulation over the contralesional motor cortex in acute stroke: a double-blind placebo controlled crossover trial. Restor Neurol Neurosci 2007;25:461–5.

Lin KC, Chang YF, Wu CY, Chen YA. Effects of constraint-induced therapy versus bilateral arm training on motor performance, daily functions, and quality of life in stroke survivors. Neurorehabil Neural Repair 2009;23:441-448.

Lin KC, Chen YA, Chen CL, Wu CY, Chang YF. The effects of bilateral arm training on motor control and functional performance in chronic stroke: a randomized controlled study. Neurorehabil Neural Repair 2010;24:42-51.

Lin KC, Wu CY, Liu JS, Chen YT, Hsu CJ. Constraint-induced therapy versus dose matched control intervention to improve motor ability, basic/extended daily functions, and quality of life in stroke. Neurorehabil Neural Repair 2009;23:160-165.

Lin KC, Wu CY, Wei TH, Lee CY, Liu JS. Effects of modified constraint-induced movement therapy on reach-to-grasp movements and functional performance after chronic stroke: a randomized controlled study. Clin Rehabil 2007;21:1075-1086.

rferg@med.umich.edu
drakoski@med.umich.edu

Lin Z, Yan T. Long-term effectiveness of neuromuscular electrical stimulation for promoting motor recovery of the upper extremity after stroke. J Rehabil Med 2011; 43(6):506-510.

Lindberg PG, Skejo PH, Rounis E, Nagy Z, Schmitz C, Wernegren H, et al. Wallerian degeneration of corticofugal tracts in chronic stroke: a pilot study relating diffusion tensor imaging, transcranial magnetic stimulation, and hand function. Neurorehabil Neural Repair 2007;21: 551–60.

Lo AC, Guarino PD, Richards LG, Haselkorn JK, Wittenberg GF, Federman DG, Ringer RJ, Wagner TH, Krebs HI, Volpe BT, Bever CT, Jr., Bravata DM, Duncan PW, Corn BH, Maffucci AD, Nadeau SE, Conroy SS, Powell JM, Huang GD, Peduzzi P. Robot-assisted therapy for long-term upper-limb impairment after stroke. N Engl J Med 2010.

Logigian MK, Samuels MA, Falconer J, Zagar R. Clinical exercise trial for stroke patients. Arch Phys Med Rehabil 1983;64(8):364-367.

Lord JP, Hall K. Neuromuscular reeducation versus traditional programs for stroke rehabilitation. Arch Phys Med Rehabil 1986;67(2):88-91.

Luft AR, McCombe-Waller S, Whitall J, et al. Repetitive bilateral arm training and motor cortex activation in chronic stroke: a randomized controlled trial. JAMA 2004;292:1853-1861.

Luke C, Dodd KJ, Brock K. Outcomes of the Bobath concept on upper limb recovery following stroke. Clin Rehabil 2004;18:888-898.

Lum PS, Burgar CG, Shor PC, Majmundar M, Van der Loos M. Robot assisted movement training compared with conventional therapy techniques for rehabilitation of upper-limb motor function after stroke. Arch Phys Med Rehabil 2002;83:952–9.

Macclellan LR, Bradham DD, Whitall J, Volpe B, Wilson PD, Ohlhoff J et al. Robotic upper limb neurorehabilitation in chronic stroke patients. J Rehabil Res Dev 2005;42(6):717-722.

Malcolm MP, Triggs WJ, Light KE, Gonzalez Rothi LJ, Wu S, Reid K, et al. Repetitive transcranial magnetic stimulation as an adjunct to constraint-induced therapy: an exploratory randomized controlled trial. Am J Phys Med Rehabil 2007;86:707–15.

Mangold S, Schuster C, Keller T, Zimmermann-Schlatter A, Ettlin T. Motor training of upper extremity with functional electrical stimulation in early stroke rehabilitation. Neurorehabil Neural Repair 2009;23:184-190.

Masiero S, Celia A, Rosita G, Armani M. Robotic-Assisted Rehabilitation of the Upper Limb After Acute Stroke. Arch Phys Med Rehabil 2007;88:142-149.

McDonnell MN, Hillier SL, Miles TS, Thompson PD, Ridding MC. Influence of combined afferent stimulation and task-specific training following stroke: a pilot randomized controlled trial. Neurorehabil Neural Repair 2007;21:435-443

rferg@med.umich.edu
drakoski@med.umich.edu

Meier KC, Oesch PR, Gamper UN, Kool JP, Beer S. Potential effectiveness of three different treatment approaches to improve minimal to moderate arm and hand function after stroke - a pilot randomized clinical trial. Clin Rehabil 2011

Merians AS, Poizner H, Boian R, Burdea G, Adamovich S. Sensorimotor Training in a Virtual Reality Environment: Does It Improve Functional Recovery Poststroke? Neurorehabil Neurol Repair 2006;20:252-267.

Michaelsen S, Dannenbaum R, Levin M. Task-specific training with trunk restraint on arm recovery in stroke. Stroke 2006;37:186–192.

Michaelsen S, Luta A, Roby-Brami A, Levin M. Effect of trunk restraint on the recovery of reaching movements in hemiparetic patients. Stroke 2001;32:1875–1883.

Michielsen ME, Selles RW, van der Geest JN, Eckhardt M, Yavuzer G, Stam HJ et al. Motor recovery and cortical reorganization after mirror therapy in chronic stroke patients: a phase II randomized controlled trial. Neurorehabil Neural Repair 2011; 25(3):223-233.

Miller GJ, Light KE. Strength training in spastic hemiparesis? Should it be avoided? NeuroRehabilitation 1997;9:17-28.

Morris JH, van WF, Joice S, Ogston SA, Cole I, MacWalter RS. A comparison of bilateral and unilateral upper-limb task training in early poststroke rehabilitation: a randomized controlled trial. Arch Phys Med Rehabil 2008;89:1237-1245.

Mouawad MR, Doust CG, Max MD, McNulty PA. Wii-based movement therapy to promote improved upper extremity function poststroke: a pilot study. J Rehabil Med 2011; 43(6):527-533.

Mudie MH, Matayas TA. Responses of the densely hemiplegic upper extremity to bilateral training. Neurorehabil Neural Repair 2001;15: 129–140.

Mudie MH, Matyas TA. Can simultaneous bilateral movement involve the undamaged hemisphere in reconstruction of neural networks damaged by stroke? Disabil Rehabil 2000;22:23-37.

Muller K, Butefisch CM, Seitz RJ, Homberg V. Mental practice improves hand function after hemiparetic stroke. Restor Neurol Neurosci 2007; 25(5-6):501-511.

Myint JM, Yuen GF, Yu TK, Kng CP,Wong AM, Chow KK, et al. A study of constraint-induced movement therapy in sub-acute stroke patients in Hong Kong. Clin Rehab 2008;22:112–24.

Nakayama H, Jorgensen HS, Raaschou HO, Olsen TS. Recovery of upper extremity function in stroke patients: the Copenhagen Stroke Study. Arch Phys Med Rehabil 1994;75:394-398.

Page SJ, Dunning K, Hermann V, Leonard A, Levine P. Longer versus shorter mental practice sessions for affected upper extremity movement after stroke: a randomized controlled trial. Clin Rehabil 2011; 25(7):627-637.

rferg@med.umich.edu
drakoski@med.umich.edu

Page SJ, Levine P, Khoury JC. Modified constraint-induced therapy combined with mental practice: thinking through better motor outcomes. Stroke 2009;40:551-554.

Page SJ, Levine P, Leonard A, Szaflarski JP, Kissela BM. Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial. Phys Ther 2008;88:333-340.

Page SJ, Levine P, Leonard A. Mental Practice in Chronic Stroke: Results of a Randomized, Placebo-Controlled Trial. Stroke 2007;38:1293-1297.

Page SJ, Levine P, Leonard AC. Effects of mental practice on affected limb use and function in chronic stroke. Arch Phys Med Rehabil 2005; 86(3):399-402.

Page SJ, Levine P, Leonard AC. Modified Constraint-Induced Therapy in Acute Stroke: A Randomized Controlled Pilot Study. Neurorehabil Neural Repair 2005;19:27-32.

Page SJ, Levine P. Back From the Brink: Electromyography-Triggered Stimulation Combined with Constraint-Induced Movement Therapy in Chronic Stroke. Arch Phys Med Rehabil 2006;87:27-31.

Page SJ. Intensity versus task-specificity after stroke: how important is intensity? Am J Phys Med Rehabil 2003;82(9):730-2.

Pang MY, Harris JE, Eng JJ. A community-based upper-extremity group exercise program improves motor function and performance of functional activities in chronic stroke: a randomized controlled trial. Arch Phys Med Rehabil 2006;87:1–9.

Piron L, Toin P, Atzori AM, Zucci Z, Massaro C, Trivello E, Dam M. The augmented feedback rehabilitation technique facilitates the arm motor recovery in patients after a recent stroke. Stud Health Technol Inform 2003;94:265-297.

Piron L, Turolla A, Agostini M, Zucconi C, Cortese F, Zampolini M, Zannini M, Dam M, Ventura L, Battauz M, Tonin P. Exercises for paretic upper limb after stroke: a combined virtual-reality and telemedicine approach. J Rehabil Med 2009;41:1016-1102.

Piron L, Turolla A, Agostini M, Zucconi CS, Ventura L, Tonin P, Dam M. Motor learning principles for rehabilitation: a pilot randomized controlled study in poststroke patients. Neurorehabil Neural Repair 2010;24:501-508.

Platz T, Eickhof C, van Kaick S, Engel U, Pinkowski C, Kalok S, et al. Impairment-oriented training or Bobath therapy for severe arm paresis after stroke: a single-blind, multicenter randomized controlled trial. Clin Rehab 2005;19:714–724.

Platz T, van KS, Mehrholz J, Leidner O, Eickhof C, Pohl M. Best conventional therapy versus modular impairment-oriented training for arm paresis after stroke: a single-blind, multicenter randomized controlled trial. Neurorehabil Neural Repair 2009;23:706-716.

rferg@med.umich.edu
drakoski@med.umich.edu

Ploughman M, Corbett D. Can forced-use therapy be clinically applied after stroke? An exploratory randomized controlled trial. Arch Phys Med Rehabil. 2004;85(9):1417-1423.

Rabadi M, Galgano M, Lynch D, Akerman M, Lesser M, Volpe B. A pilot study of activity based therapy in the arm motor recovery post stroke: a randomized controlled trial. Clin Rehabil 2008;22:1071-1082.

Rakauer JW. Motor learning: its relevance to stroke recovery and neurorehabilitation. Curr Opin Neurol 2006;19:84–90.

Renner C, Woldag H, Atanasova R, Hummelsheim H. Change of facilitation during volontary bilateral hand activation after stroke. J Neurol Sci 2005;239:25–30.

Riccio I, Iolascon G, Barillari MR, Gimigliano R, Gimigliano F. Mental practice is effective in upper limb recovery after stroke: a randomized single-blind cross-over study. Eur J Phys Rehabil Med 2010;46:19-25.

Richards L, Hanson C, Wellborn M, Sethi A, Driving motor recovery after stroke. Topics Stroke Rehabil, 2008; 15(5): 397-411.

Richards L, Rothi LJG, Davis S, Wu AA, Nadeau SE. Limited dose response to Constraint-Induced Movement therapy in patients with chronic stroke. Clin Rehabil 2006;20: 1066-1074.

Rodgers H, Mackintosh J, Price C, Wood R, McNamee P, Fearon T, Marritt A, Curless R. Does an early increased-intensity interdisciplinary upper limb therapy programme following acute stroke improve outcome? Clin Rehabil 2003;17(6):579-89.

Rose DK, Winstein CJ. The coordination of bimanual rapid aiming movements following stroke. Clin Rehab 2005;19:452–62.

Rossini PM, Rossi S. Transcranial magnetic stimulation. Diagnostic, therapeutic, and research potential. Neurology 2007;68:484–8.

Saposnik G, Teasell R, Mamdani M, Hall J, McIlroy W, Cheung D, Thorpe KE, Cohen LG, Bayley M. Effectiveness of virtual reality using Wii gaming technology in stroke rehabilitation: a pilot randomized clinical trial and proof of principle. Stroke 2010;41:1477-1484.

Sharma N, Pomeroy VM, Baron JC. Motor imagery. A backdoor to the motor system after stroke? Stroke 2006;37:1941–52.

Shumway-Cook AWM, Woollacott M. Motor control. Translating research into clinical practice. 3rd ed. Philadelphia, London: Wolters Kluwer/Lippincott Williams & Wilkins; 2007.

Stein J, Krebs HI, Frontera WR, Hogan N. Comparison of two techniques of Robot-assisted upper-limb exercise training after stroke. Am J Phys Med Rehabil 2004;83:720–728.

Stein J, Narendran K, McBean J, Krebs K, Hughes R. Electromyography-controlled exoskeletal upper-limb—powered orthosis for exercise training after stroke. Am J Phys Med Rehabil 2007;86:255-261.

rferg@med.umich.edu
drakoski@med.umich.edu

Sterr A, Szameitat A, Shen S, Freivoqel S. Application of the CIT concept in the clinical environment: hurdles, practicalities and clinical benefits. Cogn Behav Neurol 2006;19:48–54.

Stinear CM, Barber PA, Coxon JP, Fleming MK, Byblow WD. Priming the motor system enhances the effects of upper limb therapy in chronic stroke. Brain 2008;131:1381-1390.

Stinear JW, Byblow WD. Rythmic bilateral movement training modulates corticomotor excitability and enhances upper limb motricity postroke: a pilot study. J Clin Neurophysiol 2004;21:124–31.

Stoykov ME, Lewis GN, Corcos DM. Comparison of bilateral and unilateral training for upper extremity hemiparesis in stroke. Neurorehabil Neural Repair 2009;23:945-953.

Summers JJ, Kagerer FA, Garry MI, Hiraga CY, Loftus A, Cauraugh JH. Bilateral and unilateral movement training on upper limb function in chronic stroke patients: A TMS study. J Neurol Sci 2007;252:76-82.

Takeuchi N, Chuma T, Matsuo Y, Watanabe I, Ikoma K. Repetitive transcranial magnetic stimulation of contralesional primary motor cortex improves hand function after stroke. Stroke 2005;36:2681–6.

Talelli P, Greenwood RJ, Rothwell JC. Arm function after stroke: neurophysiological correlates and recovery mechanisms assessed by transcranial magnetic stimulation. Clin Neurophysiol 2006;117:1260–5.

Taub E, Lum PS, Hardin P, Mark VW, Uswatte G. AutoCITE: automated delivery of CI therapy with reduced effort by therapists. Stroke 2005;36:1301-1304.

Taub E, Miller NE, Novack TA, Cook EW, Fleming WC, Nepomuceno CS, Connell JS, Crago JE. Technique to improve chronic motor deficit after stroke. Arch Phys Med Rehabil 1993;74:347-354.

Thielman GT, Dean CM, Gentile AM. Rehabilitation of reaching after stroke: task related training versus progressive resistive exercise. Arch Phys Med Rehabil 2004;85:1613-1618.

Thrasher TA, Zivanovic V, McIlroy W, Popovic MR. Rehabilitation of reaching and grasping function in severe hemiplegic patients using functional electrical stimulation therapy. Neurorehabil Neural Repair 2008;22:706-714.

Tijs E, Matayas TA. Bilateral training does not facilitate performance of copying tasks in post-stroke hemiplegia. Neurorehabil Neural Repair 2006;20:473–83.

Trombly CA, Thayer-Nason L, Bliss G, Girard CA, Lyrist LA, Brexa-Hooson A. The effectiveness of therapy in improving finger extension in stroke patients. Am J Occup Ther 1986;40:612-617.

Underwood J, Clark PC, Blanton S, Ayick DM, Wolf SL. Pain, Fatigue and Intensity of Practice in People With Stroke Who Are Receiving Constraint-Induced Movement Therapy. Physical Therapy 2006;86:1241-1250.

van der Lee JH, Wagenaar RC, Lankhorst GJ, Vogelaar TW, Deville WL, Bouter LM. Forced use of the upper extremity in chronic stroke patients: results from a single-blind randomized clinical trial. Stroke 1999;30:2369-2375.

rferg@med.umich.edu
drakoski@med.umich.edu

van Vliet PM, Lincoln NB, Foxall A. Comparison of Bobath based and movement science based treatment for stroke: a randomized controlled trial. J Neurol Neurosurg Psychiatry 2005;76:503-508.

Volpe BT, Lynch D, Rykman-Berland A, et al. Intensive sensorimotor arm training mediated by therapist or robot improves hemiparesis in patients with chronic stroke. Neurorehabil Neural Repair 2008;22:305-310

Wagenaar RC, Meijer OG, van Wieringen PC, Kuik DJ, Hazenberg GJ, Lindeboom J, Wichers F, Rijswijk H. The functional recovery of stroke: a comparison between neuro-developmental treatment and the Brunnstrom method. Scand J Rehabil Med 1990;22(1):1-8.

Wang Q, Zhao JL, Zhu QX, Li J, Meng PP. Comparison of conventional therapy, intensive therapy and modified constraintinduced movement therapy to improve upper extremity function after stroke. J Rehabil Med 2011; 43(7):619-625.

Wang RY, Chan RC, Tsai MW. Functional electrical stimulation on chronic and acute hemiplegic shoulder subluxation. Am J Phys Med Rehabil. 2000;79(4):385-90.

Wang RY, Yang YR, Tsai MW, Wang WT, Chan RC. Effects of functional electric stimulation on upper limb motor function and shoulder range of motion in hemiplegic patients. Am J Phys Med Rehabil. 2002;81(4):283-90.

Wenzelburger R, Kopper F, Frenzel A. Hand coordination following capsular stroke. Brain 2005;128:64–74.

Whitall J, McCombe WS, Silver KH, Macko RF. Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. Stroke 2000; 31:2390-2395.

Whitall J, Waller SM, Sorkin JD, Forrester LW, Macko RF, Hanley DF et al. Bilateral and unilateral arm training improve motor function through differing neuroplastic mechanisms: a single-blinded randomized controlled trial. Neurorehabil Neural Repair 2011; 25(2):118-129.

Winstein CJ, Rose DK, Tan SM, Lewthwaite R, Chui HC, Azen SP. A randomized controlled comparison of upper-extremity rehabilitation strategies in acute stroke: A pilot study of immediate and long-term outcomes. Arch Phys Med Rehabil 2004;85:620-628.

Woldag H, Waldmann G, Heuschkel G, Hummelsheim H. Is the repetitive training of complex hand and arm movements beneficial for motor recovery in stroke patients? Clin Rehabil 2003;17:723-730.

Wolf SL, Thompson PA, Winstein CJ, Miller JP, Blanton SR, Nichols-Larsen DS, Morris DM, Uswatte G, Taub E, Light KE, Sawaki L. The EXCITE Stroke Trial. Comparing Early and Delayed Constraint-Induced Movement Therapy. Stroke 2010;41 (10): 2309-15.

Wolf SL, Winstein CJ, Miller JP, et al. Retention of upper limb function in stroke survivors who have received constraint-induced movement therapy: the EXCITE randomized trial. Lancet Neurol 2008;7:33-40.

rferg@med.umich.edu
drakoski@med.umich.edu

Wolf SL, Winstein CJ, Miller JP, Taub E, Uswatte G, Morris D, Giuliani C, Light KE, Nichols-Larsen D. Effect of Constraint-Induced Movement Therapy on Upper Extremity Function 3 to 9 months after stroke. JAMA 2006; 296: 2095-2104.

Wolf SL,Winstein C, Miller JP, Thompson PA, Taub E, Uswatte G, et al. Retention of upper limb function in stroke survivors who have received constraint-induced movement therapy: the EXCITE randomised trial. Lancet Neurol 2008;7:33–40.

Woodbury ML, Howland DR, McGuirk TE, Davis SB, Senesac CR, Kautz S, Richards LG. Effects of trunk restraint combined with intensive task practice on poststroke upper extremity reach and function: a pilot study. Neurorehabil Neural Repair 2009;23:78-91.

Wu C, Chen C, Tang S, Lin KYH. Kinematic and clinical analyses of upper-extremity movements after contraint-induced movement therapy in patients with stroke: a randomized controlled trial. Arch Phys Med Rehabil 2007;88:964–70.

Wu CY, Chuang LL, Lin KC, Chen HC, Tsay PK. Randomized trial of distributed constraint induced therapy versus bilateral arm training for the rehabilitation of upper-limb motor control and function after stroke. Neurorehabil Neural Repair 2011; 25(2):130-139.

Yavuzer G, Selles R, Sezer N, et al. Mirror therapy improves hand function in subacute stroke: a randomized controlled trial. Arch Phys Med Rehabil 2008;89:393-398.

Yavuzer G, Senel A, Atay MB, Stam HJ. "Playstation eyetoy games" improve upper extremity-related motor functioning in subacute stroke: a randomized controlled clinical trial. Eur J Phys Rehabil Med 2008;44:237-244.

Yelnik A. Evolution of the concepts concerning rehabilitation treatment for hemiplegic patients. Ann Med Phys Rehab 2005;48:270–7.

Yozbatiran N, Donmez B, Kayak N, Bozan O. Electrical stimulation of wrist and fingers for sensory and functional recovery in acute hemiplegia. Clin Rehabil 2006;20:4-11.

rferg@med.umich.edu
drakoski@med.umich.edu